






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 311/29, 311/46, 311/19, 323/67, 311/42, C07D 307/12, A61K 31/195, 31/34		A3	(11) International Publication Number: WO 98/16504
			(43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18406		(74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.	
(22) International Filing Date: 9 October 1997 (09.10.97)			
(30) Priority Data: 60/028,323 11 October 1996 (11.10.96) US		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [-/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): ALLEN, Hamish, John [GB/US]; 47 Eastern Point Drive, Shrewsbury, MA 01545 (US). BRADY, Kenneth, Dale [US/US]; 32 Ivernia Road, Worcester, MA 01606 (US). HARTER, William, Glen [US/US]; 3750 Shagbark, Chelsea, MI 48118 (US). LUNNEY, Elizabeth, Ann [US/US]; 619 Ridgewood Court, Ann Arbor, MI 48103 (US). PLUMMER, Mark, Stephen [US/US]; 8100 Huron River Drive, Dexter, MI 48130 (US). SAWYER, Tomi [US/US]; 5753 E. Silo Ridge, Ann Arbor, MI 48108 (US). SHAHRIPOUR, Aurash [US/US]; Apartment 193, 2765 Windwood Drive, Ann Arbor, MI 48105 (US). WALKER, Nigel [GB/DE]; Frauenpfad 20, D-69221 Dossenheim (DE).		Published With international search report.	
		(88) Date of publication of the international search report: 25 June 1998 (25.06.98)	

(54) Title: SULFONAMIDE SUBSTITUTED ASPARTIC ACID INTERLEUKIN-1 β CONVERTING ENZYME INHIBITORS

(57) Abstract

The present invention relates to compounds that are inhibitors of interleukin-1 β converting enzyme that have Formula (I, II or III). This invention also relates to a method of treatment of stroke, inflammatory diseases, septic shock, reperfusion injury, Alzheimer's disease, and shigellosis, and to a pharmaceutically acceptable composition that contains a compound that is an inhibitor of interleukin-1 β converting enzyme.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification ⁶ : C07C 311/29, 311/46, 311/19, 323/67, 311/42, C07D 307/12, A61K 31/195, 31/34		A3	(11) International Publication Number: WO 98/16504
			(43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18406			(74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.
(22) International Filing Date: 9 October 1997 (09.10.97)			
(30) Priority Data: 60/028,323 11 October 1996 (11.10.96) US			(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [-/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): ALLEN, Hamish, John [GB/US]; 47 Eastern Point Drive, Shrewsbury, MA 01545 (US). BRADY, Kenneth, Dale [US/US]; 32 Ivernia Road, Worcester, MA 01606 (US). HARTER, William, Glen [US/US]; 3750 Shagbark, Chelsea, MI 48118 (US). LUNNEY, Elizabeth, Ann [US/US]; 619 Ridgewood Court, Ann Arbor, MI 48103 (US). PLUMMER, Mark, Stephen [US/US]; 8100 Huron River Drive, Dexter, MI 48130 (US). SAWYER, Tomi [US/US]; 5753 E. Silo Ridge, Ann Arbor, MI 48108 (US). SHAHRIPOUR, Aurash [US/US]; Apartment 193, 2765 Windwood Drive, Ann Arbor, MI 48105 (US). WALKER, Nigel [GB/DE]; Frauenpfad 20, D-69221 Dossenheim (DE).			Published With international search report.
			(88) Date of publication of the international search report: 25 June 1998 (25.06.98)

(54) Title: SULFONAMIDE SUBSTITUTED ASPARTIC ACID INTERLEUKIN-1 β CONVERTING ENZYME INHIBITORS

(I)



(II)



(III)

(57) Abstract

The present invention relates to compounds that are inhibitors of interleukin-1 β converting enzyme that have Formula (I, II or III). This invention also relates to a method of treatment of stroke, inflammatory diseases, septic shock, reperfusion injury, Alzheimer's disease, and shigellosis, and to a pharmaceutically acceptable composition that contains a compound that is an inhibitor of interleukin-1 β converting enzyme.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/18406

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C311/29 C07C311/46 C07C311/19 C07C323/67 C07C311/42
C07D307/12 A61K31/195 A61K31/34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 496 378 A (DR. KARL THOMAE GMBH) 29 July 1992 see page 12, lines 13-24; page 14, lines 46-53; page 23, compound 41; page 27, compound 20; claims 1, 8, 9 ---	1,3-7, 10,11, 24, 32-35, 38,39, 46,47
A	WO 93 14066 A (JAMES BLACK FOUNDATION LTD) 22 July 1993 see page 1, lines 27-36; example 20; claims 1, 20 --- -/--	1,4,5, 12,34,46

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

31 March 1998

Date of mailing of the international search report

21.04.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 97/18406

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EP 0 519 748 A (MERCK & CO INC) 23 December 1992</p> <p>see page 18, lines 38-55; examples 1-3, 5, 6; claims 1, 16</p> <p>-----</p>	<p>1-4, 12, 28, 34, 36, 38, 40, 46</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 18406

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 28-45
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/18406

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 496378 A	29-07-92	DE 4102024 A	30-07-92
		AT 128120 T	15-10-95
		AU 648379 B	21-04-94
		AU 1040392 A	30-07-92
		DE 59203704 D	26-10-95
		ES 2079694 T	16-01-96
		IE 69973 B	16-10-96
		JP 4334351 A	20-11-92
		MX 9200269 A	01-07-92
		NO 177852 B	28-08-95
		NZ 241355 A	26-07-94
		US 5597825 A	28-01-97
		ZA 9200464 A	23-07-93
WO 9314066 A	22-07-93	AU 3262493 A	03-08-93
		DE 69309901 D	22-05-97
		EP 0625141 A	23-11-94
		ZA 9300137 A	08-07-94
EP 519748 A	23-12-92	CA 2071674 A	22-12-92
		JP 1971354 C	27-09-95
		JP 5255218 A	05-10-93
		JP 6102642 B	14-12-94
		US 5434248 A	18-07-95



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07C 311/00</p>	A2	<p>(11) International Publication Number: WO 98/16504</p> <p>(43) International Publication Date: 23 April 1998 (23.04.98)</p>
<p>(21) International Application Number: PCT/US97/18406</p> <p>(22) International Filing Date: 9 October 1997 (09.10.97)</p> <p>(30) Priority Data: 60/028,323 11 October 1996 (11.10.96) US</p> <p>(71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [-/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): ALLEN, Hamish, John [GB/US]; 47 Eastern Point Drive, Shrewsbury, MA 01545 (US). BRADY, Kenneth, Dale [US/US]; 32 Ivernia Road, Worcester, MA 01606 (US). HARTEK, William, Glen [US/US]; 3750 Shagbark, Chelsea, MI 48118 (US). LUNNEY, Elizabeth, Ann [US/US]; 619 Ridgewood Court, Ann Arbor, MI 48103 (US). PLUMMER, Mark, Stephen [US/US]; 8100 Huron River Drive, Dexter, MI 48130 (US). SAWYER, Tomi [US/US]; 5753 E. Silo Ridge, Ann Arbor, MI 48108 (US). SHAHRIPOUR, Aurash [US/US]; Apartment 193, 2765 Windwood Drive, Ann Arbor, MI 48105 (US). WALKER, Nigel [GB/DE]; Frauenpfad 20, D-69221 Dossenheim (DE).</p>		
<p>(74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.</p> <p>(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>		
<p>(54) Title: SULFONAMIDE SUBSTITUTED ASPARTIC ACID INTERLEUKIN-1β CONVERTING ENZYME INHIBITORS</p> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 20px;"> <div style="text-align: center;"> <p>(I)</p> </div> <div style="text-align: center;"> <p>(II)</p> </div> <div style="text-align: center;"> <p>(III)</p> </div> </div>		
<p>(57) Abstract</p> <p>The present invention relates to compounds that are inhibitors of interleukin-1β converting enzyme that have Formula (I, II or III). This invention also relates to a method of treatment of stroke, inflammatory diseases, septic shock, reperfusion injury, Alzheimer's disease, and shigellosis, and to a pharmaceutically acceptable composition that contains a compound that is an inhibitor of interleukin-1β converting enzyme.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

SULFONAMIDE SUBSTITUTED ASPARTIC ACID INTERLEUKIN-1 β
CONVERTING ENZYME INHIBITORS

FIELD OF THE INVENTION

5 This invention relates to compounds that are inhibitors of interleukin-1 β converting enzyme. This invention also relates to a method of treatment of stroke, inflammatory diseases, septic shock, reperfusion injury, Alzheimer's disease, shigellosis, and to a pharmaceutically acceptable composition that contains a compound that is an inhibitor of interleukin-1 β converting enzyme.

BACKGROUND OF THE INVENTION

10 The compounds of the present invention are inhibitors of interleukin-1 β converting enzyme (ICE) and are useful in treating diseases in which interleukin-1 plays a role.

ICE acts on pro-interleukin-1 β (pro-IL-1 β) to produce interleukin-1 β (IL-1 β), which is an inflammatory cytokine. In addition, ICE (Caspase-1) regulates
15 at least four cytokines. ICE activates IL- β and IL-18, and indirectly regulates the production of IL-1 α and IFN γ . Several diseases are associated with interleukin-1 activity. Examples of diseases in which interleukin-1 is involved include, but are not limited to, inflammatory diseases such as rheumatoid arthritis and inflammatory bowel disease, and neuroinflammatory disorders such as stroke.
20 Other diseases include septic shock, reperfusion injury, Alzheimer's disease, and shigellosis.

Agents that modulate IL-1 β activity have been shown to have beneficial in vivo effects. For example, compounds that are interleukin-1 receptor antagonists have been shown to inhibit ischaemic and excitotoxic damage in rat
25 brains. See, for example, Relton J.K., et al., Brain Research Bulletin, 1992;29:243-246. Additionally, ICE inhibitors were shown to reduce

-2-

inflammation and pyrexia in rats. See Elford P.R., et al., British Journal of Pharmacology, 1995;115:601-606.

5 The compounds of the present invention are also inhibitors of other cysteine proteases in the ICE family. Many of these proteases have only recently been described in the literature. While the nomenclature is still unresolved, the following proteases are representative members of this class of enzymes; Ich-2 (also called Tx or ICERel-II), ICERel-III, Ich-I (also called Nedd-2), CPP-32 (also called apopain and yama), Mch-2, Mch-3 (also called ICE-lap3, CMH-1), and Ced-3. See Henkart P.A., Immunity, 1996;4:195-201. It is
10 recognized that members of this enzyme family play key biological roles in both inflammation and apoptosis (programmed cell death). In particular, Caspase-4 can activate IL-1 β and IL-18. It has been shown that a murine homolog of Caspase-4 can activate ICE. Thus, inhibition of Caspase-4 will act to inhibit ICE. See Thornberry N.A., et al., Perspectives in Drug Discovery and Design,
15 1994;2:389-399.

In addition to its effects on IL-1 β production, ICE has been shown to play a role in the production of the inflammatory mediator interferon- γ (Ghayur, et al., Nature, 1997;386(6625):619-623). ICE processes the inactive proform of interferon- γ inducing factor (IGIF; Interleukin-18) to active IGIF, a protein which
20 induces production of interferon- γ by T-cells and natural killer cells. Interferon- γ has been implicated in the pathogenesis of diseases such as inflammatory disorders and septic shock. Therefore, ICE inhibitors would be expected to have beneficial effects in such disease states by effects on interferon- γ .


Recently, the nomenclature of these cysteine proteases in the ICE family
25 (also known as Caspases with ICE being known as Caspase-1) has been further defined. The following proteases are representative members of this class of enzymes using the nomenclature described in Alnemri, et al., Cell, 1996;87:171: Caspase-2 (also known as Ich-1); Caspase-3 (also known as CPP32, Yama, and apopain); Caspase-4 (also known as TX, Ich-2, and ICE rel-II); Caspase-5 (also
30 known as ICE rel-III); Caspase-6 (also known as Mch2); Caspase-7 (also known

-3-

as Mch3); Caspase-8 (also known as FLICE and Mch5); Caspase-9 (also known as ICE-LAP6 and Mch6); Caspase-10 (also known as Mch4).

SUMMARY OF THE INVENTION

Provided by the present invention are compounds having the Formula I



I

wherein

R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;

R^2 is $-\text{CHO}$, $-\text{COR}^a$, or $-\text{CN}$;

each R^a is independently hydrogen or C_1 - C_6 alkyl;

10 X is a bond, CH_2 , CHR^5 , NH , NR^5 , or O ;

R^3 is aryl, substituted-aryl, heteroaryl, substituted-heteroaryl, cycloalkyl, substituted-cycloalkyl, heterocycle, or substituted-heterocycle;

15 Y is absent, NR^5 , CO , S , O , SO_2 , $-\text{O}(\text{CHR}^5)_n$, CHR^5 , NR^5CO , NCR^5 , CONR^5 , OCHR^5 , CHR^5O , SCHR^5 , CHR^5S , SO_2NR^5 , C_1 - C_6 alkyl, NR^5SO_2 , CH_2CHR^5 , CHR^5CH_2 , COCH_2 , or CH_2CO ;

20 R^4 is absent, aryl, substituted-aryl, C_1 - C_8 alkyl, heteroaryl, substituted-heteroaryl, cycloalkyl, C_1 - C_6 alkyl, substituted-cycloalkyl, heterocycloalkyl, or substituted-heterocycloalkyl;

-4-

each R^5 is independently hydrogen, C_1 - C_6 alkyl, aryl, $-(CH_2)_n$ aryl, or $-(CH_2)_n$ cycloalkyl;

each n is independently 0 to 5, m is 1 or 2, and the pharmaceutically acceptable salts, esters, amides, and prodrugs thereof.

5 In one embodiment of the invention, R^2 is CHO.

In another embodiment of the invention, R^1 is hydrogen.

In another embodiment of the invention, R^a is hydrogen.

In another embodiment of the invention, X is a bond.

10 In another embodiment of the invention, R^3 is phenyl or substituted phenyl.

In another embodiment of the invention, Y is a bond.

In another embodiment of the invention, Y is O.


In another embodiment of the invention, Y is CH_2 .

15 In another embodiment of the invention, R^4 is phenyl or substituted phenyl.

In another embodiment of the invention, R^2 is CHO, R^a is H, R^1 is hydrogen, X is a bond, R^3 and R^4 are phenyl or substituted phenyl, and Y is a bond, CH_2 , or O.

In another embodiment of the invention, m is 1 and R^5 is hydrogen.

20 In a preferred embodiment, the present invention provides compounds of Formula II



II

-5-

wherein

R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;

R^2 is -CHO, -COR^a, or -CN;

each R^a is independently hydrogen or C_1 - C_6 alkyl;

5 X is a bond, CH_2 , CHR^5 , NH, NR^5 , or O;

Y is a bond, NR^5 , CO, S, O, SO_2 , CHR^5 , NR^5CO , $CONR^5$, $OCHR^5$, CHR^5O ,

-O(CHR^5)_n, $SCHR^5$, CHR^5S , SO_2NR^5 , NR^5SO_2 , CH_2CHR^5 ,

CHR^5CH_2 , $COCH_2$, or CH_2CO ;

each R⁵ is independently hydrogen, C_1 - C_6 alkyl, aryl, or -(CH₂)_naryl;

10 each n is independently 0 to 5;

m is 1 or 2;

Each Z is independently hydrogen, or an aryl, substituted aryl, heteroaryl,

substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocycle, or

substituted heterocycle group that is fused to the phenyl group that

15 contains Z as a substituent;

R^b, R^c, R^d, and R^e are each independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy,

-OH, C_1 - C_6 thioalkoxy, halogen, trifluoromethyl, dialkylamino, -NO₂,

-CN, -CF₃, -CO₂alkyl, -SO₃H, -CHO, -COalkyl, -CONH-alkyl,

-CONHR^q, -CON(alkyl)₂, -(CH₂)_n-NH₂, -(CH₂)_n-NH-alkyl, -NHR^q,

20 -NHCOR^q, -(CH₂)_nOH, -(CH₂)_nCONH₂, or -(CH₂)_nCO₂H; and

R^q is hydrogen or C_1 - C_6 alkyl, and the pharmaceutically acceptable salts, esters,

amides, and prodrugs thereof.

In another embodiment of the invention with respect to the compounds of

Formula II, R¹ is hydrogen.

25 In another embodiment of the invention with respect to the compounds of

Formula II, R² is CHO.

In another embodiment of the invention with respect to the compounds of

Formula II, R^a is hydrogen.

In another embodiment of the invention with respect to the compounds of Formula II, X is a bond.

In another embodiment of the invention with respect to the compounds of Formula II, Y is a bond, O, or CH₂.

5 In another embodiment of the invention with respect to the compounds of Formula II, R^b and R^c are hydrogen.

In another embodiment of the invention with respect to the compounds of Formula II, wherein R^b, R^c, and R^d are hydrogen and R^e is C₁-C₆ alkyl.

10 In another preferred embodiment of the invention with respect to the compounds of Formula II, R^b or R^c is located at the para position of the phenyl ring with respect to X and R^b or R^c is -OCH₃.

In another embodiment of the invention with respect to the compounds of Formula II, m is 1 and R⁵ is hydrogen.

15 In a more preferred embodiment, the present invention provides the compounds

3-(Biphenyl-2-sulfoamino)-4-oxo-butyric acid;

3-(2-Benzyl-benzenesulfonylamino)-4-oxo-butyric acid;

4-Oxo-3-(2-phenoxy-benzenesulfonylamino)-butyric acid;

4-Oxo-3-(2-p-tolyloxy-benzenesulfonylamino)-butyric acid;

20 3-[2-(4-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;

4-Oxo-3-(2-m-tolyloxy-benzenesulfonylamino)-butyric acid;


3-[2-(3-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;

and

3-(4'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid.

25 This invention also includes compounds of the Formula III

-7-



III

wherein

R¹ is hydrogen, C₁-C₆alkyl, or benzyl;

R² is -CHO, -COR^a, or -CN;

5 each R^a is independently hydrogen or C₁-C₆alkyl;

X is a bond, CH₂, CHR⁵, NH, NR⁵, or O;

R⁵ is hydrogen, C₁-C₆alkyl, aryl, or -(CH₂)_naryl;

each n is independently 0 to 5;

m is 1 or 2;

10 Z is absent, or an aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocycle, or substituted heterocycle group that is fused to the phenyl group that contains Z as a substituent;

R^f, R^g, are each independently hydrogen, C₁-C₆alkyl, hydroxy, halogen, trifluoromethyl, dialkylamino, -NO₂, -CN, -CO₂H, -CO₂alkyl, -SO₃H, -CHO, -COalkyl, -CONH₂, -CONH(CH₂)_naryl, -CONH(CH₂)_n-substituted-aryl, -CONH-alkyl, -CONHR^q, -CON(alkyl)₂, -(CH₂)_n-NH₂, -(CH₂)_n-NH-alkyl, -NHR^q, -NHCOR^q, -OR^q, -SR^q, or -(CH₂)_naryl; and

R^q is hydrogen or C₁-C₆alkyl, and the pharmaceutically acceptable salts, esters, amides, and prodrugs thereof.

20 In a preferred embodiment of the compounds of Formula III, R^f is ortho to X on the phenyl ring, and R^g is hydrogen.

In a preferred embodiment of the compounds of Formula III, Z is hydrogen, m is 1, R⁵ is hydrogen, and R^a is hydrogen.

In a preferred embodiment of the compounds of Formula III, the present invention provides the compound 3-benzenesulfonylamino-4-oxo-butyric acid.

5 Also provided is a method of inhibiting interleukin-1 β converting enzyme, the method comprising administering to a patient in need of inhibition of interleukin-1 β converting enzyme a therapeutically effective amount of a compound of Formula I, II, or III.

10 Also provided is a method of inhibiting Caspase-4, the method comprising administering to a patient in need of inhibition of Caspase-4 a Caspase-4 inhibiting amount of a compound of Formula I, II, or III.

 Also provided is a method of treating stroke, the method comprising administering to a patient having a stroke or having had a stroke a therapeutically effective amount of a compound of Formula I, II, or III.

15 Also provided is a method of treating inflammatory diseases, the method comprising administering to a patient having an inflammatory disease a therapeutically effective amount of a compound of Formula I, II, or III.

 Also provided is a method of treating septic shock, the method comprising administering to a patient having septic shock a therapeutically effective amount
20 of a compound of Formula I, II, or III.

 In a preferred embodiment, the inflammatory disease is arthritis or inflammatory bowel disease.

 Also provided is a method of treating reperfusion injury, the method comprising administering to a patient having reperfusion injury a therapeutically
25 effective amount of a compound of Formula I, II, or III.


 Also provided is a method of treating Alzheimer's disease, the method comprising administering to a patient having Alzheimer's disease a therapeutically effective amount of a compound of Formula I, II, or III.

30 Also provided is a method of treating shigellosis, the method comprising administering to a patient having shigellosis a therapeutically effective amount of a compound of Formulas I, II, or III.

Also provided is a pharmaceutically acceptable composition that contains a compound of Formula I, II, or III.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides compounds having the Formula I



wherein

R¹ is hydrogen, C₁-C₆alkyl, or benzyl;

R² is -CHO, -COR^a, or -CN;

each R^a is independently hydrogen or C₁-C₆alkyl;

10 X is a bond, CH₂, CHR⁵, NH, NR⁵, or O;

R³ is aryl, substituted-aryl, heteroaryl, substituted-heteroaryl, cycloalkyl, substituted-cycloalkyl, heterocycle, or substituted heterocycle;

15 Y is absent, NR⁵, CO, S, O, SO₂, -O(CHR⁵)_n, CHR⁵, NR⁵CO, NCR⁵, CONR⁵, OCHR⁵, CHR⁵O, SCHR⁵, CHR⁵S, SO₂NR⁵, C₁-C₆alkyl, NR⁵SO₂, CH₂CHR⁵, CHR⁵CH₂, COCH₂, or CH₂CO;

R⁴ is absent, aryl, substituted-aryl, C₁-C₈alkyl, heteroaryl, substituted-heteroaryl, cycloalkyl, C₁-C₆alkyl, substituted-cycloalkyl, heterocycloalkyl, or substituted heterocycloalkyl;


20

-10-

each R^5 is independently hydrogen, C_1 - C_6 alkyl, aryl, $-(CH_2)_n$ aryl, or $-(CH_2)_n$ cycloalkyl;

each n is independently 0 to 5, m is 1 or 2, and the pharmaceutically acceptable salts, esters, amides, and prodrugs thereof.

5 In a preferred embodiment, the present invention provides compounds of Formula II



wherein

R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;

10 R^2 is $-CHO$, $-COR^a$, or $-CN$;

each R^a is independently hydrogen or C_1 - C_6 alkyl;

X is a bond, CH_2 , CHR^5 , NH , NR^5 , or O ;

Y is a bond, NR^5 , CO , S , O , SO_2 , CHR^5 , NR^5CO , $CONR^5$, $OCHR^5$, CHR^5O , $-O(CHR^5)_n$, $SCHR^5$, CHR^5S , SO_2NR^5 , NR^5SO_2 , CH_2CHR^5 ,

15 CHR^5CH_2 , $COCH_2$, or CH_2CO ;

each R^5 is independently hydrogen, C_1 - C_6 alkyl, aryl, or $-(CH_2)_n$ aryl;

each n is independently 0 to 5;

m is 1 or 2;

Each Z is independently hydrogen, or an aryl, substituted aryl, heteroaryl,


20 substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocycle or substituted heterocycle group that is fused to the phenyl group that contains Z as a substituent;

-11-

R^b , R^c , R^d , and R^e are each independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, -OH, C_1 - C_6 thioalkoxy, halogen, trifluoromethyl, dialkylamino, $-NO_2$, -CN, $-CF_3$, $-CO_2$ alkyl, $-SO_3H$, -CHO, -COalkyl, -CONH-alkyl, -CONHR^q, -CON(alkyl)₂, $-(CH_2)_n-NH_2$, $-(CH_2)_n-NH$ -alkyl, -NHR^q, -NHCOR^q, $-(CH_2)_nOH$, $-(CH_2)_nCONH_2$, or $-(CH_2)_nCO_2H$; and

R^q is hydrogen or C_1 - C_6 alkyl, and the pharmaceutically acceptable salts, esters, amides, and prodrugs thereof.

The present invention also provides compounds of the Formula III



III

10 wherein

R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;

R^2 is -CHO, $-COR^a$, or -CN;

each R^a is independently hydrogen or C_1 - C_6 alkyl;

X is a bond, CH_2 , CHR^5 , NH, NR^5 , or O;

15 R^5 is hydrogen, C_1 - C_6 alkyl, aryl, or $-(CH_2)_n$ aryl;

each n is independently 0 to 5;

m is 1 or 2;

Z is absent, or an aryl, substituted aryl, heteroaryl, substituted heteroaryl,


cycloalkyl, substituted cycloalkyl, heterocycle, or substituted heterocycle

20 group that is fused to the phenyl group that contains Z as a substituent;

-12-

R^f , R^g , are each independently hydrogen, C_1 - C_6 alkyl, hydroxy, halogen, trifluoromethyl, dialkylamino, $-NO_2$, $-CN$, $-CO_2H$, $-CO_2$ alkyl, $-SO_3H$, $-CHO$, $-CO$ alkyl, $-CONH_2$, $-CONH(CH_2)_n$ aryl, $-CONH(CH_2)_n$ -substituted-aryl, $-CONH$ -alkyl, $-CONHR^q$, $-CON(alkyl)_2$, $-(CH_2)_n-NH_2$, $-(CH_2)_n-NH$ -alkyl, $-NHR^q$, $-NHCOR^q$, $-OR^q$, $-SR^q$, or $-(CH_2)_n$ aryl; and R^q is hydrogen or C_1 - C_6 alkyl, and the pharmaceutically acceptable salts, esters, amides, and prodrugs thereof.

In general, the groups X and Y in Formulas I, II, and III are spacers for the groups attached to X and Y. It is also noted that certain compounds of Formulas I, II, and III can exist in different, interconvertible forms. An example is shown below when R^1 is hydrogen and R^2 is CHO.



Both the cyclic and acyclic forms are contemplated and are considered part of the present invention. Moreover, it is preferred that the groups attached to R^3 be attached at adjacent atoms of R^3 . For example, if R^3 is phenyl, X and Y would have a 1, 2 relationship on R^3 .

The term "alkyl" means a straight or branched chain hydrocarbon. Representative examples of alkyl groups are methyl, ethyl, propyl, isopropyl, isobutyl, butyl, tert-butyl, sec-butyl, pentyl, and hexyl.

The term "alkoxy" means an alkyl group attached to an oxygen atom. Representative examples of alkoxy groups include methoxy, ethoxy, tert-butoxy, propoxy, and isobutoxy.

The term "halogen" includes chlorine, fluorine, bromine, and iodine.

The term "aryl" means an aromatic hydrocarbon. Representative examples of aryl groups include phenyl and naphthyl.

The term "heteroatom" includes oxygen, nitrogen, sulfur, and phosphorus.

The term "heteroaryl" means an aryl group wherein one or more carbon atom of the aromatic hydrocarbon has been replaced with a heteroatom. Examples of heteroaryl groups include furan, thiophene, pyrrole, thiazole, pyridine, pyrimidine, pyrazine, benzofuran, indole, coumarin, quinoline, isoquinoline, and naphthyridine.

The term "cycloalkyl" means a cyclic alkyl group. Examples of cycloalkyl groups include cyclopropane, cyclobutane, cyclopentane, and cyclohexane.

The term "heterocycle" means a cycloalkyl group on which one or more carbon atom has been replaced with a heteroatom. Examples of heterocycles include piperazine, morpholine, and piperidine.

The aryl, heteroaryl, or cycloalkyl groups may be substituted with one or more substituents, which can be the same or different. Examples of suitable substituents include alkyl, alkoxy, thioalkoxy, hydroxy, halogen, trifluoromethyl, amino, alkylamino, dialkylamino, -NO₂, -CN, -CO₂H, -CO₂alkyl, -SO₃H, -CHO, -COalkyl, -CONH₂, -CONH-alkyl, -CONHR^q, -CON(alkyl)₂, -(CH₂)_n-NH₂, -OH, -CF₃, -OC₁₋₆alkyl, -(CH₂)_n-NH-alkyl, -NHR^q, -NHCOR^q, phenyl,

O
||

-(CH₂)_nOH, -(CH₂)_nCNH₂, or -(CH₂)_nCO₂H, where n is 1 to 5 and R^q is hydrogen or alkyl.

The symbol "-" means a bond.

The compounds of Formula I, II, and III can be administered to a patient either alone or as part of a pharmaceutically acceptable composition. The compositions can be administered to patients such as humans and animals either orally, rectally, parenterally (intravenously, intramuscularly, or subcutaneously), intracisternally, intravaginally, intraperitoneally, intravesically, locally (powders, ointments, or drops), or as a buccal or nasal spray.

Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions, or emulsions, and sterile powders for reconstitution into sterile

injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles, include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid; (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia; (c) humectants, as for example, glycerol; (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate; (e) solution retarders, as for example, paraffin; (f) absorption accelerators, as for example, quaternary ammonium compounds; (g) wetting agents, as for example, cetyl alcohol and glycerol monostearate; (h) adsorbents, as for example, kaolin and bentonite; and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar, as well as, high molecular weight polyethyleneglycols, and the like.

5 Solid dosage forms such as tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They may contain opacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used are polymeric substances and waxes. The active compounds
10 can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly
15 used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofuryl alcohol,
20 polyethyleneglycols, and fatty acid esters of sorbitan or mixtures of these substances, and the like.

Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

25 Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, or mixtures of these substances, and the like.

30 Compositions for rectal administrations are preferably suppositories which can be prepared by mixing the compounds of the present invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol, or a suppository wax, which are solid at ordinary temperatures, but liquid at body

temperature and therefore, melt in the rectum or vaginal cavity and release the active component.

Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is
5 admixed under sterile conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

The compounds of the present invention can be administered to a patient at
10 dosage levels in the range of about 0.1 to about 1,000 mg per day. For a normal human adult having a body weight of about 70 kg, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is preferable. The specific dosage used, however, can vary. For example, the dosage can depend on a numbers of factors including the requirements of the patient, the severity of the
15 condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well known to those skilled in the art.

The term "pharmaceutically acceptable salts, esters, amides, and prodrugs" as used herein refers to those carboxylate salts, amino acid addition salts, esters,
20 amides, and prodrugs of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic and tautomeric forms, where possible, of the
25 compounds of the invention. The term "salts" refers to the relatively nontoxic, inorganic, and organic acid addition salts of compounds of the present invention. These salts can be prepared in situ during the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed.
30 Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate,

tartrate, naphthylate mesylate, glucoheptonate, lactobionate and laurylsulphonate salts, and the like. These may include cations based on the alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium and amine cations including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. (See, for example, Berge S.M, et al., "Pharmaceutical Salts", J. Pharm. Sci., 1977;66:1-19 which is incorporated herein by reference).

Examples of pharmaceutically acceptable, nontoxic esters of the compounds of this invention include C₁-C₆ alkyl esters wherein the alkyl group is a straight or branched chain. Acceptable esters also include C₅-C₇ cycloalkyl esters as well as arylalkyl esters such as, but not limited to benzyl. C₁-C₄ alkyl esters are preferred. Esters of the compounds of the present invention may be prepared according to conventional methods.

Examples of pharmaceutically acceptable, nontoxic amides of the compounds of this invention include amides derived from ammonia, primary C₁-C₆ alkyl amines and secondary C₁-C₆ dialkyl amines wherein the alkyl groups are straight or branched chain. In the case of secondary amines the amine may also be in the form of a 5- or 6-membered heterocycle containing one nitrogen atom. Amides derived from ammonia, C₁-C₃ alkyl primary amines, and C₁-C₂ dialkyl secondary amines are preferred. Amides of the compounds of the invention may be prepared according to conventional methods.

The term "prodrug" refers to compounds that are rapidly transformed in vivo to yield the parent compound of the above formulae, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water,

ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

The compounds of the present invention can exist in different stereoisomeric forms by virtue of the presence of asymmetric centers in the compounds; i.e., each asymmetric carbon can have either the R or S configuration. It is contemplated that all stereoisomeric forms of the compounds as well as mixtures thereof, including racemic mixtures, form part of this invention.


The compounds of the present invention are administered to a patient in need of ICE inhibition. In general, patients in need of ICE inhibition are those patients having a disease or condition in which ICE plays a role. Examples of such diseases include, but are not limited to, inflammatory diseases such as rheumatoid arthritis and inflammatory bowel disease and neuroinflammatory disorders such as stroke. Other diseases include reperfusion injury, Alzheimer's disease, and shigellosis.

A "therapeutically effective amount" is an amount of a compound of Formula I, II, or III that when administered to a patient having a disease that can be treated with a compound of Formula I ameliorates a symptom of the disease. A therapeutically effective amount of a compound of Formula I, II, or III is readily determined by one skilled in the art by administering a compound of Formula I, II, or III to a patient and observing the results.

The following examples illustrate particular embodiments of the invention and are not intended to limit the scope of the specification and claims in any manner.

The compounds of the present invention can be made generally as follows.

-19-



EXAMPLE 1

3-(Biphenyl-2-sulfoamino)-4-oxo-butyric acid

Biphenyl-2-sulfonyl chloride was obtained using a known procedure
 (Neale A.J., Rawlings T.J., McCall E.B., Tetrahedron, 1965;21:1299-1313). In
 general, the sulfonyl chloride (3.2 mmol) was stirred in dichloromethane (20 mL)
 with pyridine (12 mmol) and Asp(OtBu)-NMe(OMe) [Asp = aspartic acid;
 tBu = tert-butyl; Me = methyl] (2.9 mmol). After stirring 16 hours at room
 temperature, dichloromethane was added, and the organic layer was washed
 sequentially with 10% sulfuric acid, water, and then brine. Removal of the solvent
 provided the product as a colorless foam. Chromatography on silica gel eluting
 with 7:3 diethyl ether/hexanes gave product as a colorless foam (750 mg, 56%).
 The resulting sulfonamide was dissolved diethyl ether and cooled to -65°C .

-20-

Lithium aluminum hydride (1.5 equivalents) was added to the solution and the reaction temperature was maintained for 2 hours. The excess hydride was quenched by the addition of potassium hydrogen sulfate (2 equivalents) dissolved in water. After warming to room temperature, ether was added and the organic layer was washed with water and then brine. The solvent was evaporated to give crude product as a colorless oil. Chromatography on silica gel (3:2 ether/hexanes) gave the aldehyde (327 mg, 50%). The aldehyde was treated with 3:1 dichloromethane/trifluoroacetic acid (20 mL) containing a trace of water for 2 hours. The solvent was evaporated, excess trifluoroacetic acid was chased with toluene and then ether to give product as a colorless foam (242 mg, 90%).
¹H NMR (CD₃OD) as the lactol 8.05 (m, 1H), 7.60 (t, 1H), 7.53 (t, 1H), 7.41 (m, 5H), 7.30 (d, 1H), 4.42 (s, 1H), 3.63 (s, 1H), 2.50 (m, 1H), 2.30 (m, 1H);
Mass Spectrometry - Chemical Ionization (MS CI) + 1% NH₃ in CH₄ 334 (M+H)⁺.

EXAMPLE 2

3-Benzenesulfonylamino-4-oxo-butyric acid

Benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam employing the methods previously described in Example 1. ¹H NMR (CD₃OD) as the lactol 7.87 (m, 2H), 7.58 (m, 3H), 4.44, 4.38 (d, 1H), 3.74 (m, 1H), 2.57 (m, 1H), 2.27 (m, 1H);
MS CI + 1% NH₃ in CH₄ 240 (M-OH)⁺.

EXAMPLE 3

3-(2-Benzyl-benzenesulfonylamino)-4-oxo-butyric acid

Diphenylmethane-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam (62 mg, 9%) employing the methods previously described in Example 1. ¹H NMR (CD₃OD) as the lactol 8.00 (dd, 1H), 7.42 (t,

-21-

1H), 7.25 (m, 6H), 7.05 (dd, 1H), 4.54, 4.50 (d, 1H), 4.01 (s, 2H), 3.63 (m, 1H), 2.58 (m, 1H), 2.37 (m, 1H);

MS CI + 1% NH₃ in CH₄ 304 (M-CHO)⁺.

EXAMPLE 4

5 4-Oxo-3-(2-phenoxy-benzenesulfonylamino)-butyric acid

Diphenylether-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam (72 mg, 7%) employing the methods previously described in Example 1. ¹H NMR (CD₃OD) as the lactol 7.95 (d, 1H), 7.43 (m, 10 3H), 7.20 (m, 4H), 6.82 (d, 1H), 4.58, 4.52 (d, 1H), 3.75 (m, 1H), 2.70-2.40 (m, 2H);

MS APCI probe temperature 450°C, cone voltage 25 volts, acetonitrile: methanol 4:1, 350.5 (M+H)⁺.

EXAMPLE 5

15 4-Oxo-3-(2-p-tolyloxy-benzenesulfonylamino)-butyric acid

(4'-Methylphenyl) phenylether-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam (152 mg, 15%) employing the methods previously described in Example 1. ¹H NMR (CD₃OD) as the lactol 7.90 (d, 1H), 20 7.48 (d, 1H), 7.35-7.00 (m, 5H), 6.80 (d, 1H), 4.55, 4.50 (d, 1H), 3.75 (m, 1H), 2.70-2.42 (m, 2H), 2.35 (s, 3H);

MS APCI probe temperature. 450°C, cone voltage 25 volts, acetonitrile: methanol 4:1, 363.1 (M+H)⁺.

EXAMPLE 6

25 3-[2-(4-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid

(4'-Isopropylphenyl)phenylether-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam (248 mg, 22%) employing the methods

previously described in Example 1. ^1H NMR (CD_3OD) as the lactol 7.92 (d, 1H), 7.50 (t, 1H), 7.32 (m, 2H), 7.13 (m, 3H), 6.82 (d, 1H), 4.58, 4.52 (d, 1H), 3.75 (m, 1H), 2.95 (m, 1H), 2.70-2.42 (m, 2H), 1.25 (d, 6H);

MS APCI probe temperature 450°C , cone voltage 25 volts, acetonitrile: methanol

5 4:1, 392.4 ($\text{M}+\text{H}$) $^+$.

EXAMPLE 7

4-Oxo-3-(2-m-tolyloxy-benzenesulfonylamino)-butyric acid

(3'-Methylphenyl) phenylether-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final
10 product was obtained as a colorless foam (85 mg, 8%) employing the methods previously described in Example 1. ^1H NMR (CD_3OD) as the lactol 7.90 (d, 1H), 7.50 (t, 1H), 7.35-7.13 (m, 2H), 7.10-6.80 (m, 4H), 4.58-4.52 (d, 1H), 3.76 (m, 1H), 2.65-2.42 (m, 2H), 2.35 (s, 3H);

MS APCI probe temperature 450°C , cone voltage 25 volts, acetonitrile: methanol

15 4:1, 363.1 ($\text{M}+\text{H}$) $^+$.

EXAMPLE 8

3-[2-(3-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid

(3'-Isopropylphenyl) phenylether-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final
20 product was obtained as a colorless foam (334 mg, 29%) employing the methods previously described in Example 1. ^1H NMR (CD_3OD) as the lactol 7.90 (d, 1H), 7.50 (t, 1H), 7.45 (m, 1H), 7.25-6.80 (m, 5H), 4.58, 4.52 (d, 1H), 3.78 (m, 1H), 2.90 (m, 1H), 2.70-2.40 (m, 2H), 1.25 (d, 6H);

MS APCI probe temperature 450°C , cone voltage 25 volts, acetonitrile: methanol

25 4:1, 392.5 ($\text{M}+\text{H}$) $^+$.

EXAMPLE 9

3-(4'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid

(4'-Methyl)biphenyl-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe); from the resulting sulfonamide, final product was obtained as a colorless foam (334 mg, 29%) employing the methods previously described.

^1H NMR (CD_3OD) as the lactol 8.03 (m, 1H), 7.60 (m, 1H), 7.50 (m, 1H), 7.35 (m, 3H), 7.21 (m, 2H), 4.41 (d, 1H), 4.60 (m, 1H), 2.45 (m, 1H), 2.40 (s, 3H), 2.35 (m, 1H);

MS APCI probe temperature 450°C , cone voltage 25 volts, acetonitrile: methanol 4:1, 348.5 (M+H) $^+$.

EXAMPLE 10

3-(2-Isobutoxy-benzenesulfonylamino)-4-oxo-butyric acid

2-(Isobutoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (112 mg, 56%) employing methods previously described. ^1H NMR (CD_3OD) as the lactol 7.81 (m, 1H), 7.58 (m, 1H), 7.17 (dd, 1H), 7.05 (m, 1H), 4.50, 4.40 (d, 1H), 3.85 (d, 2H), 3.61 (m, 1H), 2.45 (m, 2H), 2.20 (m, 1H), 1.04 (d, 6H); MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile, 330.5 (M+H) $^+$.

EXAMPLE 11

3-[2-(2-Methyl-pentanoylamino)-benzenesulfonylamino]-4-oxo-butyric acid

2-Nitrobenzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe) employing methods previously described. The resulting nitrosulfonamide was reduced employing 5%Pd on carbon and hydrogen gas to provide the aminosulfonamide ^1H NMR (CDCl_3) 7.70 (d, 1H), 7.30 (t, 1H), 6.76 (m, 2H), 5.92 (d, 1H), 4.60 (m, 1H), 3.63 (s, 3H), 3.02 (s, 3H), 2.62-2.39 (m, 2H), 1.42 (s, 9H); MS CI + 1% NH_3 in CH_4 332 (M-tBu) $^+$. The aminosulfonamide

-24-

(400 mg, 1.0 mmol) was acylated employing 2-methylpentanoyl chloride (1.5 mmol) and triethylamine (3 mmol) in dichloromethane (5 mL).

Chromatography of the crude reaction product (4:1, diethyl ether, hexane) gave the acylated amine (148 mg, 30%) ^1H NMR (CDCl_3) 9.34 (d, 1H), 8.55 (dd, 1H),

5 7.86 (d, 1H), 7.55 (t, 1H), 7.17 (t, 1H), 5.90 (d, 1H), 4.51 (m, 1H), 3.58 (s, 3H), 3.03 (m, 3H), 2.43 (m, 3H), 1.80-1.25 (m, 4H), 1.40 (s, 9H), 1.22 (d, 3H), 0.94 (t, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts,

acetonitrile:methanol 4:1, 486.6 ($\text{M}+\text{H}$) $^+$. Final product was obtained from this sulfonamide employing methods previously described. ^1H NMR (CD_3OD) as the
10 lactol 8.30 (m, 1H), 7.93 (m, 1H), 7.64-7.20 (m, 4H), 4.50-4.25 (m, 2H), 3.63 (m, 1H), 2.73-2.20 (m, 3H), 1.65 (m, 1H), 1.42 (m, 2H), 1.22 (d, 3H), 0.97 (t, 3H).

EXAMPLE 12

4-Oxo-3-(2-o-tolyloxy-benzenesulfonylamino)-butyric acid

2-(o-Tolyloxy)-benzenesulfonyl chloride was employed to acylate the
15 amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (206 mg, 66%) employing methods previously described. ^1H NMR (CD_3OD) as the lactol; 7.90 (m, 1H), 7.43 (m, 1H), 7.38-7.08 (m, 5H), 6.62 (d, 1H), 4.58 (dd, 1H), 3.79 (m, 1H), 2.70-2.42 (m, 2H), 2.03 (m, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts,
20 acetonitrile:methanol 4:1, 364.4 ($\text{M}+\text{H}$) $^+$.

EXAMPLE 13

4-Oxo-3-(2-phenethyl-benzenesulfonylamino)-butyric acid

2-(Phenethyl)-benzenesulfonyl chloride was employed to acylate the amine
of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was
25 obtained as a colorless foam (56 mg, 25%) employing methods previously described. ^1H NMR (CD_3OD) as the lactol; 8.00 (m, 1H), 7.60-7.10 (m, 8H) 4.50, 4.45 (d, 1H), 3.63 (m, 1H), 3.30 (m, 2H), 2.97 (m, 2H), 2.62-2.42 (m, 2H);

-25-

MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 362.5 (M+H)⁺.

EXAMPLE 14

3-(2-Cyclohexyloxy-benzenesulfonylamino)-4-oxo-butyric acid

2-(Cyclohexyloxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (190 mg, 95%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.80 (m, 1H), 7.57 (m, 1H), 7.23-6.98 (m, 2H), 4.58 (m, 1H), 4.50, 4.40 (d, 1H), 3.63 (m, 1H), 2.45 (m, 2H), 2.10-1.40 (m, 10H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 356.5 (M+H)⁺.

EXAMPLE 15

3-[2-(1-Chloro-naphthalen-2-yloxy)-benzenesulfonylamino]-4-oxo-butyric acid

2-(1-Chloro-naphthalen-2-yloxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (199 mg, 92%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 8.30 (d, 1H), 7.93 (m, 3H), 7.70 (m, 1H), 7.60 (m, 1H), 7.48 (m, 1H), 7.39 (m, 1H), 7.20 (m, 1H), 6.75 (d, 1H), 4.63, 4.58 (d, 1H), 3.83 (m, 1H), 2.40 (m, 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 434.4 (M+H)⁺.

EXAMPLE 16

4-Oxo-3-[2-(5, 6, 7, 8-tetrahydro-naphthalen-2-yloxy)-benzenesulfonylamino]-butyric acid

2-(5,6,7,8-tetrahydro-naphthalen-2-yloxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (205 mg, 87%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.90 (m, 1H), 7.42 (m, 1H), 7.17 (m, 2H), 7.00 (d, 2H), 6.90 (m, 1H), 6.62 (d, 2H),

-26-

4.58, 4.46 (d, 1H), 3.78 (m, 1H), 2.81 (m, 2H), 2.63 (m, 2H), 2.50 (m, 2H), 1.80 (m, 4H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 404.5 (M+H)⁺.

EXAMPLE 17

5 4-Oxo-3-(2-phenethyloxy-benzenesulfonylamino)-butyric acid

2-phenethyloxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (224 mg, 69%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.78 (m, 1H), 7.55 (m, 1H), 7.42-7.00 (m, 7H), 4.35 (m, 2H), 4.30, 4.20 (d, 1H), 3.55 (m, 1H), 3.20 (t, 2H), 2.52-2.14 (m, 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 378.4 (M+H)⁺.

EXAMPLE 18

15 3-[2-(2-Ethyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyrlic acid

2-(2-Ethyl-phenoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (295 mg, 70%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.92 (m, 1H), 7.40 (m, 2H), 7.30-7.05 (m, 4H), 6.63 (d, 1H), 4.58, 4.56 (d, 1H), 3.78 (m, 1H), 2.60-2.42 (m, 4H), 1.16 (t, 1H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 378.4 (M+H)⁺.

EXAMPLE 19

25 3-[2-(4-sec-Butyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyrlic acid

2-(4-sec-Butyl-phenoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (295 mg, 70%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.92 (m, 1H), 7.48 (t, 1H), 7.30-7.07 (m, sH), 6.82 (d, 1H), 4.56, 4.53 (d, 1H), 3.75 (m, 1H), 2.70-2.43 (m,

-27-

3H), 1.62 (m, 2H), 1.24 (d, 3H), 0.83 (t, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 406.5 (M+H)⁺.

EXAMPLE 20

3-[2-(Biphenyl-4-yloxy)-benzenesulfonylamino]-4-oxo-butyric acid

5 2-(Biphenyl-4-yloxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (334 mg, 88%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.95 (m, 1H), 7.63 (m, 4H), 7.58-7.08 (m, 7H), 6.94 (d, 1H), 4.59, 4.55 (d, 1H), 3.75 (m, 1H), 2.57 (m, 10 2H); electrospray MS (50:50 acetonitrile:water +0.1% NH₄OH) m/z 424.1 (M-H)⁻.

EXAMPLE 21

3-{2-[4-(1-Methyl-pentyl)-phenoxy]-benzenesulfonylamino}-4-oxo-butyric acid

15 2-[4-(1-Methyl-pentyl)-phenoxy]-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (158 mg, 69%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.90 (m, 1H), 7.47 (m, 1H), 7.30-7.05 (m, 5H), 6.82 (d, 1H), 4.58, 4.52 (d, 1H), 3.75 (m, 1H), 2.80-2.43 (m, 3H), 1.58 (q, 2H), 1.12 (m, 7H), 0.83 (t, 3H); electrospray MS 20 (50:50 acetonitrile:water +0.1% NH₄OH) m/z 432.1 (M-H)⁻.

EXAMPLE 22

3-[2-(4-Isopropyl-3-methyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid

25 2-(4-Isopropyl-3-methyl-phenoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (382 mg, 79%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.87 (m, 1H), 7.45 (t, 1H), 7.30 (m, 1H), 7.15 (m, 1H), 6.95 (m, 2H), 6.82 (d, 1H), 4.58, 4.52 (d, 1H), 3.75

(m, 1H), 3.17 (m, 1H), 2.57 (m, 2H), 2.33 (s, 3H), 1.23 (d, 6H); electrospray MS (50:50 acetonitrile:water +0.1% NH₄OH) m/z 404.0.1 (M-H)⁻.

EXAMPLE 23

3-(2-Benzoyloxy-benzenesulfonylamino)-4-oxo-butyric acid

5 2-Benzoyloxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (137 mg, 91%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.83 (m, 1H), 7.55 (m, 3H), 7.37 (m, 3H), 7.20 (m, 1H), 7.06 (m, 1H), 5.28 (s, 2H), 4.45, 4.41 (d, 1H), 3.64 (m, 10 1H)2.60-2.30 (m, 2H), MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 362.0 (M-H)⁻.

EXAMPLE 24

3-[2-(2, 3-Dimethyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid

15 2-(2,3-Dimethyl-phenoxy) -benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (373 mg, 82%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.90 (m, 1H), 7.42 (m, 1H), 7.17 (m, 3H), 6.97 (m, 1H), 6.58 (d, 1H), 4.58, 4.55 (d, 1H), 3.79 (m, 1H), 2.68-2.44 (m, 2H), 2.33 (s, 3H), 2.14 (s, 3H)MS APCI Probe temperature 450°C, 20 cone voltage 25 volts, acetonitrile:methanol 4:1, 375.8 (M-H)⁻.

EXAMPLE 25

3-{2-[4-(1-Ethyl-propyl)-phenoxy]-benzenesulfonylamino}-4-oxo-butyric acid

25 2-[4-(1-Ethyl-propyl)-phenoxy] -benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (467 mg, 67%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.92 (m, 1H), 7.45 (t, 1H), 7.17 (m, 5H), 6.83 (d, 1H), 4.58, 4.53 (d, 1H), 3.77 (m, 1H), 2.68-2.30 (m, 3H),

1.73 (m, 2H), 1.58 (m, 2H), 0.79 (t, 6H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 417.9 (M-H)⁻.

EXAMPLE 26

4-Oxo-3-[2-(3, 4, 5-trimethyl-phenoxy)-benzenesulfonylamino]-butyric acid

5 2-(3,4,5-trimethyl-phenoxy) -benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (407 mg, 79%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.88 (m, 1H), 7.44 (t, 1H), 7.16 (m, 2H), 6.80 (m, 3H), 4.58, 4.54 (d, 1H), 3.75 (m, 1H), 2.65-2.43 (m, 2H),
10 2.28 (s, 6H), 2.18 (s, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 389.8 (M-H)⁻.

EXAMPLE 27

3-(2-sec-Butoxy-benzenesulfonylamino)-4-oxo-butyrlic acid

15 2-sec-Butoxy -benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (450 mg, 73%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.80 (m, 1H), 7.55 (m, 1H), 7.18 (m, 1H), 7.02 (m, 1H), 4.70 (m, 1H), 4.53, 4.42 (d, 1H), 3.65 (m, 1H), 2.45 (m, 2H), 1.88 (m, 1H), 1.75-1.38 (m, 3H), 1.38 (dd, 3H), 0.97 (dt, 3H); MS APCI Probe
20 temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 342.1 (M-H)⁻.

EXAMPLE 28

3-[2-(4-sec-Butyl-3-methyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyrlic acid

25 2-(4-sec-Butyl-3-methyl-phenoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (257 mg, 69%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.90m, 1H), 7.45 (t, 1H), 7.18 (m, 2H), 6.98 (m, 2H), 6.83 (d, 1H), 4.57, 4.53 (d, 1H), 3.75 (m, 1H), 2.94

-30-

(m, 1H), 2.58 (m, 2H), 2.32 (s, 3H), 1.63 (m, 2H), 1.21 (d, 3H), 0.86 (t, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 420.1 (M+H)⁺.

EXAMPLE 29

5 3-(2-Ethylbutoxy-benzenesulfonylamino)-4-oxo-butyric acid

2-Ethylbutoxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (113 mg, 51%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.85 (m, 1H), 7.65 (m, 1H), 7.17 (m, 10 1H), 7.00 (m, 1H), 4.45, 4.35 (d, 1H), 4.05 (m, 2H), 3.72 (m, 1H), 2.45 (m, 2H), 1.77 (m, 1H), 1.58 (m, 4H), 0.95 (t, 6H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 356.1 (M-H)⁻.

EXAMPLE 30

15 3-(2-Methylbutoxy-benzenesulfonylamino)-4-oxo-butyric acid

2-Methylbutoxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (450 mg, 73%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.80 (m, 1H), 7.55 (m, 1H), 7.18 (m, 1H), 7.04 (m, 1H), 4.68 (m, 1H), 4.52, 4.42 (d, 1H), 3.65 (m, 1H), 2.43 (m, 2H), 20 1.84 (m, 1H), 1.75-1.40 (m, 3H), 1.37 (m, 3H), MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 342.1 (M-H)⁻.

EXAMPLE 31

25 4-Oxo-3-[2-(1-naphthyl)ethyloxy]-benzenesulfonylamino)-butyric acid

2-(1-Naphthyl)ethyloxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a pink foam (120 mg, 90%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 8.23 (d, 1H), 7.92 (d, 1H), 7.80 (m, 2H), 7.55 (m, 4H), 7.15 (m, 2H), 7.03 (m, 1H), 4.55 (m, 2H), 4.20, 4.03

-31-

(d, 1H), 3.74 (m, 2H), 3.60 (m, 1H), 2.44-2.10 (m, 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 428.1 (M+H)⁺.

EXAMPLE 32

5 3-(2'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid

2'-Methyl-biphenyl-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (70 mg, 84%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol 8.08 (m, 1H), 7.61 (m, 1H), 7.58 (m, 10 1H), 7.10-7.25 (m, 5H), 4.52, 4.41 (d, 1H), 3.70 (m, 1H), 2.45 (m, 2H), 2.03 (s, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile, 348.4 (M+H)⁺.

EXAMPLE 33

15 3-(2-Naphthalen-1-yl-benzenesulfonylamino)-4-oxo-butyric acid

2-Naphthalen-1-yl-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (110 mg, 71%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 8.18 (m, 1H), 7.90 (m, 2H), 7.65 (m, 2H), 7.58-7.40 (m, 3H), 7.30 (m, 3H), 4.42, 4.40 (d, 1H), 3.63 (m, 1H), 2.40 (m, 20 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 384.4 (M+H)⁺.

EXAMPLE 34

25 3-(Naphthalene-1-sulfonylamino)-4-oxo-butyric acid

Naphthalene-1-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (90 mg, 66%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 8.85 (m, 1H), 8.25 (t, 1H), 8.16 (t,

-32-

1H), 8.00 (d, 1H) 7.72-7.52 (m, 3H), 4.38, 4.09 (dd, 1H), 3.67 (m, 1H), 2.80-2.22 (m, 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 308.5 (M+H)⁺.

EXAMPLE 35

5 3-(3'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid

3'-Methyl-biphenyl-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (160 mg, 76%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 8.07 (m, 1H), 7.60 (m, 1H), 7.52 (m, 10 1H), 7.38-7.08 (m, 5H), 4.43 (m, 1H), 2.43 (m, 1H), 2.38 (s, 3H), 2.25 (m, 1H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 348.3 (M+H)⁺.

EXAMPLE 36

15 3-[2-(Naphthalen-2-yloxy)-benzenesulfonylamino]-4-oxo-butyric acid

2-(Naphthalen-2-yloxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (60 mg, 37%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.98 (m, 2H), 7.87 (m, 1H), 7.81 (m, 1H), 7.60 (m, 1H), 7.45 (m, 3H), 7.38 (m, 1H), 7.20 (m, 1H), 6.90 20 (m, 1H), 4.60, 4.57 (d, 1H), 3.80 (m, 1H), 2.65-2.44 (m, 2H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 399.8 (M+H)⁺.

EXAMPLE 37

25 3-(6-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid

6-Methyl-biphenyl-2-sulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (120 mg, 86%) employing methods previously

described. ^1H NMR (CD_3OD) as the lactol; 7.92 (m, 1H), 7.52 (d, 1H), 7.40 (m, 4H), 7.30 (t, 1H), 7.20 (t, 1H), 4.46 (d, 1H), 3.65 (m, 1H), 2.52-2.32 (m, 2H), 1.98 (s, 3H); MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:methanol 4:1, 348.10 ($\text{M}+\text{H}$) $^+$.

5

EXAMPLE 38

3-(3-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid

2-(Phenyloxy)-3-methylbenzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (100 mg, 55%) employing methods previously described. ^1H NMR (CD_3OD) as the lactol; 7.82 (t, 1H), 7.53 (t, 1H), 7.32 (m, 1H), 7.25 (t, 2H), 7.02 (t, 1H), 6.81 (d, 2H), 4.52, 4.42 (d, 1H), 3.75 (m, 1H), 2.54 (m, 1H), 2.42 (m, 1H), 2.04 (s, 3H); MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:methanol 4:1, 361.9 ($\text{M}-\text{H}$) $^-$.

10

EXAMPLE 39

15 4-Oxo-3-[2-phenylthio-benzenesulfonylamino]-butyric acid

2-Phenylthio-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (140 mg, 80%) employing methods previously described. ^1H NMR (CD_3OD) as the lactol; 7.95 (m, 1H), 7.69 (m, 1H), 7.44 (m, 2H), 7.35 (t, 1H), 7.27 (t, 1H), 7.00 (d, 1H), 4.56, 4.53 (d, 1H), 3.75 (m, 1H), 2.60 (m, 1H), 2.44 (m, 1H); MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:methanol 4:1, 363.8 ($\text{M}-\text{H}$) $^-$.

20

EXAMPLE 40

4-Oxo-3-[2-N-phenyl-benzenesulfonylamino]-butyric acid

2-N-phenyl-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (25 mg, 18%) employing methods previously

25

described. ¹H NMR (CD₃OD) as the lactol; 7.81 (m, 1H), 7.36 (m, 3H), 7.24 (m, 3H), 7.15 (t, 1H), 6.98 (t, 1H), 6.88 (t, 1H), 4.52, 4.45 (d, 1H), 3.65 (m, 1H), 2.62 (m, 1H), 2.38 (m, 1H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 347.0 (M-H)⁻.

5

EXAMPLE 41

3-[2-(4-Isopropylphenoxy)-3-methylbenzenesulfonylamino]-4-oxo-butyrac acid

4-Isopropylphenyloxy-3-methylbenzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (110 mg, 82%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.82 (t, 1H), 7.53 (t, 1H), 7.30 (m, 1H), 7.13 (d, 2H), 6.73 (d, 2H), 4.52, 4.42 (d, 1H), 3.74 (q, 1H), 2.58 (m, 1H), 2.42 (m, 1H), 2.03 (s, 3H), 1.20 (s, 6H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 406.1 (M+H)⁺.

10

EXAMPLE 42

15

3-[2-(2-Methylphenoxy)-3-methylbenzenesulfonylamino]-4-oxo-butyrac acid

2-Methylphenyloxy-3-methylbenzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (150 mg, 66%) employing methods previously described. ¹H NMR (CD₃OD) as the lactol; 7.80 (m, 1H), 7.51 (m, 1H), 7.32 (m, 1H), 7.10 (t, 1H), 6.98 (t, 1H), 6.86 (m, 1H), 6.28 (m, 1H), 4.42, 4.28 (d, 1H), 3.68 (m, 1H), 2.58 (m, 1H), 2.44 (s, 3H), 2.38 (m, 1H), 1.95 (s, 3H); MS APCI Probe temperature 450°C, cone voltage 25 volts, acetonitrile:methanol 4:1, 378.1 (M+H)⁺.

20

EXAMPLE 43

25

4-Oxo-3-[2-(tetrahydro-furan-2-ylmethoxy)-benzenesulfonylamino]-butyrac acid

2-(Tetrahydro-furan-2-ylmethoxy)-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a thick oil (0.1033 g, 15%) employing methods

previously described. ^1H NMR (CD_3OD , ppm) as the lactol; 7.9-7.7 (m, 1H), 7.7-7.5 (m, 1H), 7.3-7.0 (m, 2H), 4.5-4.2 (m, 3H), 4.2-3.6 (m, 4H), 2.7-2.2 (m, 2H), 2.2-1.7 (m, 4H). MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:methanol 4:1, 358.1 (M+H) $^+$.

5

EXAMPLE 44

3-(5-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid

5-Methyl-2-phenoxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a pink foam (0.2274 g, 30%) employing methods

10

previously described. ^1H NMR (CD_3OD , ppm) as the lactol; 7.73 (m, 1H), 7.5-7.0 (m, 7H), 6.76 (d, 1H), 4.5 (dd, 1H), 3.7-3.6 (m, 1H), 2.7-2.4 (m, 2H), 2.36 (s, 3H). MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:methanol 4:1, 362.9 (M-H) $^-$.

EXAMPLE 45

15

3-(4-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid

4-Methyl-2-phenoxy-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a colorless foam (0.1279 g, 15%) employing methods previously described. ^1H NMR (CD_3OD , ppm) as the lactol; 7.8 (m, 1H), 7.4 (m, 2H), 7.3-6.9 (m, 4H), 6.65 (s, 1H), 4.5 (dd, 1H), 3.7 (m, 1H), 2.7-2.4 (m, 2H), 2.28 (s, 3H). MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:water 4:1, 364.1 (M+H) $^+$.

20

EXAMPLE 46

3-(2-Nitro-benzenesulfonylamino)-4-oxo-butyric acid

25

2-Nitro-benzenesulfonyl chloride was employed to acylate the amine of Asp(OtBu)-NMe(OMe). From the resulting sulfonamide, final product was obtained as a thick yellow oil (0.028 g, 4%) employing methods previously

described. ^1H NMR (CD_3OD , ppm) as the lactol; 8.11 (m, 1H), 7.88 (m, 1H), 7.79 (m, 2H), 4.50 (m, 1H), 3.83 (m, 1H), 2.7-2.4 (m, 2H). MS APCI Probe temperature 450°C , cone voltage 25 volts, acetonitrile:water 4:1, 301.9 (M) $^-$.

INHIBITION STUDIES

5 Compounds of Formulas I, II, or III are inhibitors of ICE as demonstrated by measurement of K_i (μM) and IC_{50} (μM) using the protocol described herein. ICE (0.24 nM final concentration) is added to 400 μL of HGDE buffer (100 mM HEPES, 20% glycerol, 5 mM DTT, 0.5 mM EDTA) containing 15 μM substrate (Ac-Tyr-Val-Ala-Asp-AMC; $K_M = 15 \mu\text{M}$) plus vehicle (DMSO) or inhibitor at
10 concentrations bracketing the K_i . Substrate hydrolysis is monitored for 300 seconds by observing the fluorescence of released AMC using excitation at 380 nm and emission at 460 nm. Mean rates of substrate hydrolysis are evaluated by linear-regression analysis of the fluorescence vs time traces. To evaluate K_i , plots of percent inhibition vs inhibitor concentration are fit by non-linear
15 regression to a reversible, competitive model:

$$\% \text{Inhibition} = \frac{100 * [I]}{[I] + K_i * \left(1 + \frac{[S]}{K_M} \right)}$$

where the competition factor $(1 + [S]/K_M) = 2$.

ICE Colorimetric Dose-Response (IC_{50}) Assay

20 Diluted inhibitor stocks are prepared by two-fold serial dilution from a primary stock whose concentration is selected (based on screening results or on prior attempts at IC_{50} evaluation) to achieve approximately 95% inhibition in the most concentrated well. Aliquots of each dilution are transferred to a microtitre plate in triplicate.

ICE enzyme is diluted to approximately 24 nM in HGE buffer (100 mM Hepes pH 7.5, 0.5 mM EDTA, 20% glycerol, 0.1% Bovine Serum Albumin (BSA), and activated by adding dithiothreitol (DTT) to a final concentration of 5 mM. The activated enzyme is then aliquoted into wells containing inhibitor or vehicle, and the plate is preincubated for 60 minutes at ambient temperature. Substrate (Ac-Tyr-Val-Ala-Asp-pNA) is added to each well to a final concentration of 50 μ M, and plates are placed in the microtitre plate-reader thermostated to 25°C. Beginning 5 minutes after addition of substrate, absorbance (405 nm) of wells is monitored for 1 hour, and activity is calculated as the mean rate of change in absorbance during this interval.

PBMC Cellular Assay (IC₅₀) Determinations

Further evidence that compounds of Formula I, II, and III are inhibitors of ICE is provided by their ability to inhibit IL-1 β production in human peripheral blood mononuclear cells (PBMCs) as described herein. PBMCs are isolated from heparinized blood by centrifugation over a ficoll cushion, then washed three times with phosphate-buffered saline. PBMCs are suspended in a medium containing RPMI 1640 with glutamine, penicillin, streptomycin, and 2% human AB serum, then plated at 10⁶ cells per well in 96 well flat bottom plates. PBMCs are stimulated overnight with 10 ng/mL of lipopolysaccharide (LPS, E. Coli strain 0111:B4; Calbiochem) in the presence or absence of a compound of Formula I. Medium is harvested and the level of mature IL-1 β was determined using an ELISA kit from R & D Systems. Compound inhibition is assessed by determining the concentration of agent which reduces IL-1 β levels by 50%. Cells were cultured for an additional four hours in the presence of 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) to determine viability. Compound toxicity can, therefore, be assessed by determining the concentration of agent which kills 50% of the cells (IC₅₀).

Ich-2 (Caspase-4) Colorimetric Dose-Response (IC₅₀) Assay

Inhibition of Ich-2 enzyme is assayed as described above for ICE, except that enzyme is used at 64 nM, and 60 μ M of the Ich-2-specific substrate Ac-Leu-Glu-Val-Asp-pNA is used instead of the ICE substrate Ac-Tyr-Val-Ala-Asp-pNA.

5 The results of these test are shown below in Table 1.

TABLE 1

Example Number	ICE K _i (μ M)	ICE IC ₅₀ (μ M)	PBMC IC ₅₀ (μ M)	PBMC TC ₅₀ (μ M)	Ich-2 (Caspase-4) IC ₅₀ (μ M)
1	1.6	20.3	>100	>100	88
2	6.8	86.8	>100	ND	259
3	3.7	37.0	ND	ND	70
4	2.4	24.0	ND	ND	52
5	1.1	8.8	67.5	>100	18
6	0.54	5.3	23.5	>100	6.1
7	0.43	10.8	67.5	>100	33
8	0.41	3.7	45.0	>100	6.1
9	6.4	66.2	ND	ND	109
10	4.2	22	90	>100	318
11	17	55	ND	ND	569
12	3	14	90	>100	59
13	7	32	ND	ND	167
14	2.3	20	90	>100	35
15	1	5.1	50	>100	22
16	1.3	5.8	65	>100	42
17	0.9	9.2	55	>100	63
18	2	14	80	>100	87
19	0.5	2.2	23	>100	4.0
20	0.3	4.4	40	>100	17
21	0.6	6.1	33	70	31
22	0.11	1.0	25	>100	76
23	0.8	11	60	>100	97
24	1.0	13	73	>100	122
25	0.44	3.5	27	>100	6.5
26	0.45	7.8	58	>100	23

TABLE 1

Example Number	ICE K_i (μ M)	ICE IC_{50} (μ M)	PBMC IC_{50} (μ M)	PBMC TC_{50} (μ M)	Ich-2 (Caspase-4) IC_{50} (μ M)
27	3.9	35	>100	>100	118
28	0.1	1.0	18	>100	4.6
29	0.3	8.6	75	>100	215
30	4.2	37	ND	ND	221
31	0.062	0.40	8.0	>100	2.3
32	7	28	ND	ND	603
33	7	11.4	ND	ND	77
34	31	110	ND	ND	602
35	31	37	>100	>100	341
36	0.18	3.5	50	>100	28
37	1.9	34	>100	>100	151
38	0.67	11.2	85	>100	76
39	1.1	21	>100	>100	25
40	10	44	95	>100	370
41	1.5	12	45	>100	9.2
42	0.30	1.1	38	>100	57
43	7.7	99	ND	ND	1,973
44	2.2	41	>100	>100	199
45	18	1.7	70	>100	14
46	10	130	ND	ND	525

ND = Not determined

HEPES = 4-(2-hydroxymethyl)-1-piperazine ethane sulfonic acid

DTT = Dithiothreitol

EDTA = Ethylene diamine tetra acetic acid

AMC = 7-amino-4-methyl coumarin

Tyr = Tyrosine

Val = Valine

Ala = Alanine

Asp = Aspartic Acid

pNA = Para nitroaniline

LEU = Leucine

Glu = Glutamic acid


tBu = tert-butyl

Me = Methyl

APCI = Atmospheric pressure chemical ionization

CLAIMS

1. A compound having the Formula I



wherein

- 5 R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;
 R^2 is -CHO, -COR^a, or -CN;
each R^a is independently hydrogen or C_1 - C_6 alkyl;
X is a bond, CH_2 , CHR^5 , NH, NR^5 , or O;
 R^3 is aryl, substituted-aryl, heteroaryl, substituted-heteroaryl, cycloalkyl,
10 substituted-cycloalkyl, heterocycle, or substituted heterocycle;
Y is absent, NR^5 , CO, S, O, SO_2 , $-O(CHR^5)_n$, CHR^5 ,
 NR^5CO , $-NCR^5$, $CONR^5$, $OCHR^5$, CHR^5O , $SCHR^5$, CHR^5S ,
15 SO_2NR^5 , C_1 - C_6 alkyl, NR^5SO_2 , CH_2CHR^5 , CHR^5CH_2 , $COCH_2$,
or CH_2CO ;
 R^4 is absent, aryl, substituted-aryl, C_1 - C_8 alkyl, heteroaryl, substituted-
heteroaryl, cycloalkyl, C_1 - C_6 alkyl, substituted-cycloalkyl,
heterocycloalkyl, or substituted heterocycloalkyl;
20 each R^5 is independently hydrogen, C_1 - C_6 alkyl, aryl, $-(CH_2)_n$ aryl, or
 $-(CH_2)_n$ cycloalkyl;


-41-

each n is independently 0 to 5, m is 1 or 2, and the pharmaceutically acceptable salts, esters, amides and prodrugs thereof.

2. A compound according to Claim 1 wherein R^2 is CHO.
3. A compound according to Claim 1 wherein R^1 is hydrogen.
- 5 4. A compound according to Claim 1 wherein R^a is hydrogen.
5. A compound according to Claim 1 wherein X is a bond.
6. A compound according to Claim 1 wherein R^3 is phenyl or substituted phenyl.
7. A compound according to Claim 1 wherein Y is a bond.
- 10 8. A compound according to Claim 1 wherein Y is O.
9. A compound according to Claim 1 wherein Y is CH_2 .
10. A compound according to Claim 1 wherein R^4 is phenyl or substituted phenyl.
11. A compound according to Claim 1 wherein R^2 is CHO, R^a is H, R^1 is
15 hydrogen, X is a bond, R^3 and R^4 are phenyl or substituted phenyl, and Y is a bond, CH_2 , or O.
12. A compound according to Claim 1 wherein m is 1 and R^5 is hydrogen.

-42-

13. A compound of Formula II



wherein

 R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;5 R^2 is -CHO, -COR^a, or -CN;each R^a is independently hydrogen or C_1 - C_6 alkyl; X is a bond, CH_2 , CHR^5 , NH , NR^5 , or O ; Y is a bond, NR^5 , CO , S , O , SO_2 , CHR^5 , NR^5CO , $CONR^5$, $OCHR^5$,- $O(CHR^5)_n$ -, CHR^5O , $SCHR^5$, CHR^5S , SO_2NR^5 , NR^5SO_2 ,10 CH_2CHR^5 , CHR^5CH_2 , $COCH_2$, or CH_2CO ;each R^5 is independently hydrogen, C_1 - C_6 alkyl, aryl, or $-(CH_2)_n$ aryl;each n is independently 0 to 5; m is 1 or 2;Each Z is independently hydrogen, or an aryl, substituted aryl, heteroaryl,

15 substituted heteroaryl, cycloalkyl, substituted cycloalkyl,

heterocycle, or substituted heterocycle group that is fused to the

phenyl group that contains Z as a substituent; R^b , R^c , R^d , and R^e are each independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, -OH, C_1 - C_6 thioalkoxy, halogen, trifluoromethyl,20 dialkylamino, -NO₂, -CN, -CF₃, -CO₂alkyl, -SO₃H, -CHO,-COalkyl, -CONH-alkyl, -CONHR^q, -CON(alkyl)₂, $-(CH_2)_n$ -NH₂,

-43-

$-(CH_2)_n-NH-alkyl$, $-NHR^q$, $-NHCOR^q$, $-(CH_2)_nOH$,

$-(CH_2)_nCONH_2$, or $-(CH_2)_nCO_2H$; and


R^q is hydrogen or C_1-C_6 alkyl, and the pharmaceutically acceptable salts, esters, amides and prodrugs thereof.

- 5 14. A compound according to Claim 13 wherein R^1 is hydrogen.
15. A compound according to Claim 13 wherein R^2 is CHO.
16. A compound according to Claim 13 wherein R^a is hydrogen.
17. A compound according to Claim 13 wherein X is a bond.
18. A compound according to Claim 13 wherein Y is a bond, O, or CH_2 .
- 10 19. A compound according to Claim 13 wherein R^b and R^c are hydrogen.
20. A compound according to Claim 13 wherein R^b , R^c , and R^d are hydrogen and R^e is C_1-C_6 alkyl.
21. A compound according to Claim 13 wherein R^b or R^c is located at the para position of the phenyl ring with respect to X and R^b or R^c is $-OCH_3$.
- 15 22. A compound according to Claim 13 wherein is 1 and R^5 is hydrogen.
23. The compounds
 - 3-(Biphenyl-2-sulfoamino)-4-oxo-butyric acid;
 - 3-(2-Benzyl-benzenesulfonylamino)-4-oxo-butyric acid;
 - 4-Oxo-3-(2-phenoxy-benzenesulfonylamino)-butyric acid;
 - 20 4-Oxo-3-(2-p-tolyloxy-benzenesulfonylamino)-butyric acid;

-44-

3-[2-(4-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;
 4-Oxo-3-(2-m-tolyloxy-benzenesulfonylamino)-butyric acid;
 3-[2-(3-Isopropyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid; or
 3-(4'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid.

5 24. A compound of the Formula III



wherein

R^1 is hydrogen, C_1 - C_6 alkyl, or benzyl;

R^2 is -CHO, -COR^a, or -CN;

10 each R^a is independently hydrogen or C_1 - C_6 alkyl;

X is a bond, CH_2 , CHR^5 , NH , NR^5 , or O ;

R^5 is hydrogen, C_1 - C_6 alkyl, aryl, or $-(CH_2)_n$ aryl;

each n is independently 0 to 5;

m is 1 or 2;

15 Z is absent, or an aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocycle, or substituted heterocycle group that is fused to the phenyl group that contains Z as a substituent;

20 R^f , R^g , are each independently hydrogen, C_1 - C_6 alkyl, hydroxy, halogen, trifluoromethyl, dialkylamino, -NO₂, -CN, -CO₂H, -CO₂alkyl, -SO₃H, -CHO, -COalkyl, -CONH₂, -CONH(CH₂)_naryl,

-45-

-CONH(CH₂)_n-substituted-aryl, -CONH-alkyl, -CONHR^q,
-CON(alkyl)₂, -(CH₂)_n-NH₂, -(CH₂)_n-NH-alkyl, -NHR^q,
-NHCOR^q, -OR^q, -SR^q, or -(CH₂)_naryl; and

R^q is hydrogen or C₁-C₈alkyl, and the pharmaceutically acceptable salts,
5 esters, amides, and prodrugs thereof.

25. A compound in accordance with Claim 24 wherein R^f is ortho to X on the
phenyl ring and R^g is hydrogen.

26. A compound in accordance with Claim 24 wherein Z is hydrogen, m is 1,
R⁵ is hydrogen, and R⁹ is hydrogen.

10 27. The compound 3-benzenesulfonylamino-4-oxo-butyric acid.

28. A method of inhibiting interleukin-1 β converting enzyme, the method
comprising administering to a patient in need of inhibition of
interleukin-1 β converting enzyme a therapeutically effective amount of a
compound of Claim 1.

15 29. A method of inhibiting interleukin-1 β converting enzyme, the method
comprising administering to a patient in need of inhibition of
interleukin-1 β converting enzyme a therapeutically effective amount of a
compound of Claim 24.

20 30. A method of inhibiting Caspase-4, the method comprising administering to
a patient in need of Caspase-4 inhibition a Caspase-4 inhibiting amount of
a compound of Claim 1.

31. A method of inhibiting Caspase-4, the method comprising administering to
a patient in need of Caspase-4 inhibition a Caspase-4 inhibiting amount of
a compound of Claim 24.

32. A method of treating stroke, the method comprising administering to a patient having a stroke or having had a stroke a therapeutically effective amount of a compound of Claim 1.
- 5 33. A method of treating stroke, the method comprising administering to a patient having a stroke or having had a stroke a therapeutically effective amount of a compound of Claim 24.
34. A method of treating inflammatory diseases, the method comprising administering to a patient having an inflammatory disease a therapeutically effective amount of a compound of Claim 1.
- 10 35. A method of treating inflammatory diseases, the method comprising administering to a patient having an inflammatory disease a therapeutically effective amount of a compound of Claim 24.
36. The method of Claim 35 wherein the inflammatory disease is arthritis or inflammatory bowel disease.
- 15 37. The method of Claim 35 wherein the inflammatory disease is arthritis or inflammatory bowel disease.
38. A method of treating septic shock, the method comprising administering to a patient having septic shock a therapeutically effective amount of a compound of Claim 1.
- 20 39. A method of treating septic shock, the method comprising administering to a patient having septic shock a therapeutically effective amount of a compound of Claim 24.
- 25 40. A method of treating reperfusion injury, the method of comprising administering to a patient having reperfusion injury a therapeutically effective amount of a compound of Claim 1.

-47-

41. A method of treating reperfusion injury, the method of comprising administering to a patient having reperfusion injury a therapeutically effective amount of a compound of Claim 24.
- 5 42. A method of treating Alzheimer's disease, the method comprising administering to a patient having Alzheimer's disease a therapeutically effective amount of a compound of Claim 1.
43. A method of treating Alzheimer's disease, the method comprising administering to a patient having Alzheimer's disease a therapeutically effective amount of a compound of Claim 24.
- 10 44. A method of treating shigellosis, the method comprising administering to a patient having shigellosis a therapeutically effective amount of a compound of Claim 1.
45. A method of treating shigellosis, the method comprising administering to a patient having shigellosis a therapeutically effective amount of a compound of Claim 24.
- 15 46. A pharmaceutically acceptable composition that contains a compound of Claim 1.
47. A pharmaceutically acceptable composition that contains a compound of Claim 24.
- 20 48. The compounds:
3-(2-Isobutoxy-benzenesulfonylamino)-4-oxo-butyric acid;
3-[2-(2-Methyl-pentanoylamino)-benzenesulfonylamino]-4-oxo-butyric acid;
4-Oxo-3-(2-o-tolyloxy-benzenesulfonylamino)-butyric acid;
25 4-Oxo-3-(2-phenethyl-benzenesulfonylamino)-butyric acid;

- 3-(2-Cyclohexyloxy-benzenesulfonylamino)-4-oxo-butyric acid;
3-[2-(1-Chloro-naphthalen-2-yloxy)-benzenesulfonylamino]-4-oxo-butyric
acid;
4-Oxo-3-[2-(5, 6, 7, 8-tetrahydro-naphthalen-2-yloxy)-
5 benzenesulfonylamino]-butyric acid;
4-Oxo-3-(2-phenethyloxy-benzenesulfonylamino)-butyric acid;
3-[2-(2-Ethyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;
3-[2-(4-sec-Butyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;
3-[2-(Biphenyl-4-yloxy)-benzenesulfonylamino]-4-oxo-butyric acid;
10 3-{2-[4-(1-Methyl-pentyl)-phenoxy]-benzenesulfonylamino}-4-oxo-
butyric acid;
3-[2-(4-Isopropyl-3-methyl-phenoxy)-benzenesulfonylamino]-4-oxo-
butyric acid;
3-(2-Benzoyloxy-benzenesulfonylamino)-4-oxo-butyric acid;
15 3-[2-(2, 3-Dimethyl-phenoxy)-benzenesulfonylamino]-4-oxo-butyric acid;
3-{2-[4-(1-Ethyl-propyl)-phenoxy]-benzenesulfonylamino}-4-oxo-butyric
acid;
4-Oxo-3-[2-(3, 4, 5-trimethyl-phenoxy)-benzenesulfonylamino]-butyric
acid;
20 3-(2-sec-Butoxy-benzenesulfonylamino)-4-oxo-butyric acid;
3-[2-(4-sec-Butyl-3-methyl-phenoxy)-benzenesulfonylamino]-4-oxo-
butyric acid;
3-(2-Ethylbutoxy-benzenesulfonylamino)-4-oxo-butyric acid;
3-(2-Methylbutoxy-benzenesulfonylamino)-4-oxo-butyric acid;
25 4-Oxo-3-[2-(1-naphthyl)ethyloxy]-benzenesulfonylamino)-butyric acid;
3-(2'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid;
3-(2-Naphthalen-1-yl-benzenesulfonylamino)-4-oxo-butyric acid;
3-(Naphthalene-1-sulfonylamino)-4-oxo-butyric acid;
3-(3'-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid;
30 3-[2-(Naphthalen-2-yloxy)-benzenesulfonylamino]-4-oxo-butyric acid;
3-(6-Methyl-biphenyl-2-sulfonylamino)-4-oxo-butyric acid;

- 3-(3-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid;
4-Oxo-3-[2-phenylthio-benzenesulfonylamino]-butyric acid;
4-Oxo-3-[2-N-phenyl-benzenesulfonylamino]-butyric acid;
3-[2-(4-Isopropylphenoxy)-3-methylbenzenesulfonylamino]-4-oxo-butyric
5 acid;
3-[2-(2-Methylphenoxy)-3-methylbenzenesulfonylamino]-4-oxo-butyric
acid;
4-Oxo-3-[2-(tetrahydro-furan-2-ylmethoxy)-benzenesulfonylamino]-
butyric acid;
10 3-(5-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid;
3-(4-Methyl-2-phenoxy-benzenesulfonylamino)-4-oxo-butyric acid; and
3-(2-Nitro-benzenesulfonylamino)-4-oxo-butyric acid.